

International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB10797 - Goat Anti-MOB2 (aa128-140) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: MOB2, HCCA2, HCCA2 protein, mob2 homolog, mps one binder kinase activator-like 2

Official Symbol: MOB2

Accession Number(s): NP_001165694.1; NP_443731.2

Human GeneID(s): [81532](#)

Non-Human GeneID(s): 101513 (mouse), 499288 (rat)

Important Comments: This antibody is expected to recognize both reported isoforms (NP_001165694.1; NP_443731.2) only.

Immunogen

Peptide with sequence DERGKKVKCTAPQ, from the internal region of the protein sequence according to NP_001165694.1; NP_443731.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 30kDa band observed in lysates of cell line Jurkat (calculated MW of 30.2kDa according to NP_001165694.1). Recommended concentration: 0.02-0.06µg/ml. An additional band of unknown identity was also consistently observed at 48kDa. This band was successfully blocked by incubation with the immunizing peptide.

IHC: In paraffin embedded Human Lung shows staining of smooth muscle cells, vasculature and apical secretory granules in the epithelium of the bronchiole.

Recommended concentration: 5-10µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

Specific Reference


This antibody has been successfully used in Western blot on Rat:

Fang KM, Liu YY, Lin CH, Fan SS, Tsai CH, Tzeng SF.


Mps one binder 2 gene upregulation in the stellation of astrocytes induced by cAMP-dependent pathway.

J Cell Biochem. 2012 Sep;113(9):3019-28. doi: 10.1002/jcb.24180.

PMID: 22566124



EB10797 (0.02 μ g/ml) staining of Jurkat lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.



EB10797 (5 μ g/ml) staining of paraffin embedded Human Lung. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.