

International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB10288 - Goat Anti-MOGAT2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: acyl CoA:monoacylglycerol acyltransferase 2, DGAT2L5, FLJ22644, MGAT2, MGC119183, MGC119184, MGC119185, monoacylglycerol O-acyltransferase 2, MOGAT2

Official Symbol: MOGAT2

Accession Number(s): NP_079374.2

Human GeneID(s): [80168](#)

Immunogen

Peptide with sequence C-KESAAHILNRK, from the internal region of the protein sequence according to NP_079374.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Approx 37kDa band observed in Human Colon, Duodenum and Ileum lysates (calculated MW of 38.2kDa according to NP_079374.2). Recommended concentration: 0.05-0.2µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Small Intestine. Recommended concentration: 3.75µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog, Cow, Pig



EB10288 (0.05µg/ml) staining of Human Duodenum lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB10288 (3.75µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer Ph 6, AP-staining.