

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08705 - Goat Anti-MRP8 / ABCC11 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** ABCC11, ATP-binding cassette transporter MRP8, ATP-binding cassette, sub-family C (CFTR/MRP), member 11, EWWD, MRP8, WW, ATP-binding cassette protein C11, ATP-binding cassette, sub-family C, member 11, multi-resistance

protein 8

Official Symbol: ABCC11

Accession Number(s): NP\_115972.2; NP\_660187.1; NP\_001357425.1

Human GeneID(s): 85320

### **Immunogen**

Peptide with sequence C-KVVEFDRPEVLRK, from the internal region (near C Terminus) of the protein sequence according to NP\_115972.2; NP\_660187.1; NP\_001357425.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 125kDa band observed in Human Testis lysates and in preliminary testing of Breast lysate (calculated MW of 150kDa according to NP\_660187.1). This molecular weight is routinely observed by other sources. Recommended concentration: 0.5-1.5µg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Breast. Recommended concentration: 3.75µg/ml.

**Immunofluorescence:** Strong expression of the protein seen in the membranes and cytoplasm of HeLa cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of HeLa cells. Recommended concentration: 10ug/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa

50kDa


37kDa

25kDa


20kDa

15kDa


EB08705 (0.5 $\mu$ g/ml) staining of Human Testes lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.



 $EB08705~(3.75 \mu g/ml)~staining~of~paraffin~embedded~Human~Breast.~Steamed~antigen~retrieval~with~citrate~buffer~pH~6,~AP-staining.$ 



EB08705 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB08705 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control:

Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.