

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08489 - Goat Anti-MTHFD1 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** MTHFD1, methylenetetrahydrofolate dehydrogenase (NADP+ dependent) 1, methenyltetrahydrofolate cyclohydrolase, formyltetrahydrofolate synthetase, MTHFC, MTHFD, 5,10-methylenetetrahydrofolate dehydrogenase, 5,10-methylenetetrahydrofolate cyclohydrolase, 10-formyltetrahydrofolate synthetase, C1-THF synthase, NADP-dependent cyclohydrolase/formyltetrahydrofolate synthetase, cytoplasmic C-1-tetrahydrofolate synthase, methylenetetrahydrofolate dehydrogenase (NADP+ dependent), methenyltetrahydrofolate cyclohydrolase, formyltetrahydrofolate synthetase, methylenetetrahydrofolate dehydrogenase 1

Official Symbol: MTHFD1

Accession Number(s): NP\_005947.2; NP\_001351766.1

Human GeneID(s): 4522

Non-Human GeneID(s): 108156 (mouse), 64300 (rat)

### **Immunogen**

Peptide with sequence RGDLNDCFIPCTPK, from the internal region of the protein sequence according to NP\_005947.2; NP\_001351766.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx. 100kDa band observed in lysates of cell lines Jurkat and MOLT4 and approx.110kDa in lysates of cell line MCF7 (calculated MW of 102kDa according to NP\_005947.2). Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Liver. Recommended concentration: 3.75µg/ml.

**Immunofluorescence:** Strong expression of the protein seen in the cytoplasm of MCF7 cells. Recommended concentration:  $10\mu g/ml$ .


### **Species Reactivity**

Tested: Human


Expected from sequence similarity: Human, Mouse, Rat



EB08489 (1μg/ml) staining of Jurkat (A), MCF7 (B) and (2ug/ml) MOLT4 (C) cel lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB08489 (3.75 $\mu$ g/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB08489 Immunofluorescence analysis of paraformaldehyde fixed MCF7 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).