

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10910 - Goat Anti-MTHFD1L (aa535-538) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: MTHFD1L, methylenetetrahydrofolate dehydrogenase (NADP+ dependent) 1-like, DKFZp586G1517, FLJ21145, FTHFSDC1, MTC1THFS, dJ292B18.2, 10-formyl-THF synthetase, formyltetrahydrofolate synthetase domain containing 1

Official Symbol: MTHFD1L

Accession Number(s): NP_056255.2

Human GenelD(s): 25902

Non-Human GeneID(s): 270685 (mouse), 361472 (rat)

Immunogen

Peptide with sequence C-DTNDRFLRKITIGQ, from the internal region of the protein sequence according to NP_056255.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 110kDa band observed in Human Brain (Cerebellum) (calculated MW of 106kDa according to NP_056255.2). The same band is observed in lysates of cell line NIH3T3, Mouse Brain and Rat Brain. Recommended concentration: 0.1-0.3μg/ml. Primary incubation was 1 hour. Preliminary testing was unsuccessful on Pig for this particular batch

IHC: Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.

Additional validation: This antibody has been successfully used in the following paper: Sikorski et al. (2018) PMID: 30377371.

Species Reactivity

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

Specific Reference

This antibody has been successfully used in the following paper:

Krzysztof Sikorski, Adi Mehta, Marit Inngjerdingen, Flourina Thakor, Simon Kling, Tomas Kalina, Tuula A. Nyman, Maria Ekman Stensland, Wei Zhou, Gustavo A. De Souza, Lars Holden, Jan Stuchly, Markus Templin and Fridtjof Lund-Johansen


A high-throughput pipeline for validation of antibodies

Nat Methods. 2018 Nov;15(11):909-912


PMID: 30377371

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB10910 (0.1μg/ml) staining of Human Cerebellum lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB10910 (0.1 μ g/ml) staining of NIH3T3 (A), Mouse Brain (B) and Rat Brain (C) lysates (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB10910 (5μg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.