



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11385 - Goat Anti-MTOR / FRAP2 (aa1137-1150) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: FK506 binding protein 12-rapamycin associated protein 1, FK506 binding protein 12-rapamycin associated protein 2, FK506-binding protein 12-rapamycin complex-associated protein 1, FKBP12-rapamycin complex-associated protein 1, FKBP-rapamycin associated protein, FLJ44809, FRAP, FRAP1, FRAP2, mammalian target of rapamycin, mechanistic target of rapamycin (serine/threonine kinase), RAFT1, rapamycin and FKBP12 target 1, rapamycin associated protein FRAP2, rapamycin target protein 1, RAPT1, serine/threonine-protein kinase mTOR, MTOR

Official Symbol: MTOR

Accession Number(s): NP_004949.1

Human GenelD(s): [2475](#)

Non-Human GenelD(s): 56717 (mouse), 56718 (rat)

Immunogen

Peptide with sequence C-ETVDRLTESLDFTD, from the internal region of the protein sequence according to NP_004949.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:64000.

Immunofluorescence: Strong expression of the protein seen in the ER and vesicles/cytoplasm of HEK293 cells. Recommended concentration: 10µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB11385 Immunofluorescence analysis of paraformaldehyde fixe HEK293 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing endoplasmic reticulum and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).