

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08371 - Goat Anti-MUC13 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: MUC13, mucin 13, cell surface associated, DRCC1, FLJ20063,

down-regulated in colon cancer 1, mucin 13, epithelial transmembrane

Official Symbol: MUC13

Accession Number(s): NP\_149038.3

Human GenelD(s): 56667

## **Immunogen**

Peptide with sequence DCKSDLQRPNPQ, from the internal region of the protein sequence according to NP\_149038.3.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:4000.

**Immunofluorescence:** Strong expression of the protein seen in the cytoplasm of HepG2 and Caco-2 cells. Recommended concentration: 10µg/ml.


# **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB08371 Immunofluorescence analysis of paraformaldehyde fixed HepG2 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB08371 Immunofluorescence analysis of paraformaldehyde fixed Caco-2 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).