

#### International Office

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB10629 - Goat Anti-Muc19 / Smgc Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** Smgc, submandibular gland protein C, 2310010P21Rik, DXImx49e, Muc19, Sfc21, OTTMUSP00000022832, neonatal submandibular gland protein C

Official Symbol: Smgc

Accession Number(s): NP\_945121.1 Non-Human GeneID(s): 223809 (mouse)

#### **Immunogen**

Peptide with sequence C-KLEPKYENPTNGS, from the internal region of the protein sequence according to NP\_945121.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.


Western blot: Preliminary experiments gave bands at approx 65kDa and 70kDa in Mouse fetal Lung lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the bands we observe given the calculated size of 74.4kDa according to NP\_945121.1. Both detected bands were successfully blocked by incubation with the immunizing peptide (and BLAST results with the immunizing peptide sequence did not identify any other proteins to explain the additional bands). We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

**Immunocytochemistry:** Positive staining of select epithelial cells in the murine Submanibular gland at E18. Recommended concentration 4ug/ml. Data provided by Everest Grant winner Melinda Larsen, State University of New York, Albany, NY.

## **Species Reactivity**

Tested: Mouse

Expected from sequence similarity: Mouse



EB10629 (4ug/ml) staining of Mouse SubMandibular Gland cells at E18. Nuclear counterstain with DAPI in green.