



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB10658 - Goat Anti-Muc5b (mouse) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: Muc5b, mucin 5, subtype B, tracheobronchial, 2300002I04Rik, A130042M24, AV085033, MUC5, MUC9, mucin 5b

Official Symbol: Muc5b

Accession Number(s): NP_083077.2

Human GeneID(s): [727897](#)

Non-Human GeneID(s): 74180 (mouse), 309114 (rat)

Important Comments: EB06890

Immunogen

Peptide with sequence QHTYTHIDEEN, from the internal region of the protein sequence according to NP_083077.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.


Western blot: Not yet tested - our routine western blotting protocol does not allow for the detection of proteins >250kDa.

Immunofluorescence: Strong expression of the protein seen in vesicles of NIH3T3 cells. Recommended concentration: 10µg/ml.

Species Reactivity

Tested: Mouse

Expected from sequence similarity: Mouse, Rat



EB10658 Immunofluorescence analysis of paraformaldehyde fixed NIH3T3 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).