



UK Office

Everest Biotech Ltd
Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06680 - Goat Anti-NANOS1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: NANOS1, nanos homolog 1 (Drosophila), NOS1, nanos homolog 1

Official Symbol: NANOS1

Accession Number(s): NP_955631.1

Human GeneID(s): [340719](#)

Immunogen

Peptide with sequence C-SARDGPPGKKLR, from the C Terminus of the protein sequence according to NP_955631.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 35-37kDa band observed in Human Testis lysates (calculated MW of 30.2kDa according to NP_955631.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Clarke et al, Hum Mol Genet. 2004 Apr 1;13(7):727-39. Epub 2004 Feb 12, PMID: 14962983). Recommended concentration: 1-3µg/ml. Primary incubation was 1 hour. This antibody has been successfully used in WB on Human: Sajek et al (2018) PMID: 30269240.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human

Specific Reference


This antibody has been successfully used in Western blot on Human:

Sajek M, Janecki DM, Smialek MJ, Ginter-Matuszewska B, Spik A, Oczkowski S, Ilaslan E, Kusz-Zamelczyk K, Kotecki M, Blazewicz J, Jaruzelska J.

PUM1 and PUM2 exhibit different modes of regulation for SIAH1 that involve cooperativity with NANOS paralogues

Cell Mol Life Sci. 2018 Sep 29.

PMID: 30269240



EB06680 (0.3 μ g/ml) staining of Human Testis lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.