



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06991 - Goat Anti-NCAM2 / OCAM Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** NCAM2, neural cell adhesion molecule 2, HGNC:7657, MGC51008, NCAM21, OCAM

**Official Symbol:** NCAM2

**Accession Number(s):** NP\_004531.2

**Human GeneID(s):** [4685](#)

**Non-Human GeneID(s):** 17968 (mouse), 288280 (rat)

### Immunogen

Peptide with sequence C-KVSNDIQSKEDDSK, from the C Terminus of the protein sequence according to NP\_004531.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:4000.

**Western blot:** This antibody has been successfully used in WB on Mouse in the following papers, PMID: 34299022 and PMID: 32043120.

**IHC:** Frozen section of Olfactory bulb from transcardially PFA-perfused Mouse shows strong staining of axon bundles (right side middle, figure 1) and moderate peripheral staining outside the glomeruli (top, figure 1). NCAM2 (red) is stained complementary to VGLUT2 (blue) and NQO1 (green) in figure 2. Recommended concentration: 0.5µg/ml. Paraffin embedded Human Brain (Cortex). Recommended concentration: 3.75ug/ml. This antibody has been successfully used in IHC in the following papers, PMID: 37724425 and PMID: 32043120.

### Species Reactivity

**Tested:** Human, Mouse

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow

### Specific References

**This antibody has been successfully used in IHC:**

Alba Ortega-Gascó, Antoni Parcerisas, Keiko Hino, Vicente Herranz-Pérez, Fausto Ulloa, Alba Elias-Tersa, Miquel Bosch, José Manuel García-Verdugo, Sergi Simó, Lluís Pujadas, Eduardo Soriano

Regulation of young-adult neurogenesis and neuronal differentiation by neural cell adhesion molecule 2 (NCAM2).

Cereb Cortex. 2023 Oct 14;33(21):10931-10948.

PMID: 37724425

**This antibody has been successfully used in WB on Mouse:**

Antoni Parcerisas, Alba Ortega-Gascó, Marc Hernaiz-Llorens, Maria Antonia Odena, Fausto Ulloa, Eliandre de Oliveira, Miquel Bosch, Lluís Pujadas, and Eduardo

Soriano

New Partners Identified by Mass Spectrometry Assay Reveal Functions of NCAM2 in Neural Cytoskeleton Organization.

Int J Mol Sci. 2021 Jul; 22(14): 7404.

PMID: 34299022


**This antibody has been successfully used in WB and IHC on Mouse:**

Antoni Parcerisas, Lluís Pujadas, Alba Ortega-Gascó, Bartomeu Perelló-Amorós, Ricardo Viais, Keiko Hino, Joana Figueiro-Silva, Anna La Torre, Ramón Trullás, Sergi Simó, Jens Lüders, and Eduardo Soriano


NCAM2 Regulates Dendritic and Axonal Differentiation through the Cytoskeletal Proteins MAP2 and 14-3-3.

Cereb Cortex. 2020 May; 30(6): 3781-3799.


PMID: 32043120



EB06991 (0.5 $\mu$ g/ml) 24h-staining of PFA-perfused cryosection of Mouse Olfactory bub. Antigen retrieval with methanol (-20C, 10min) followed by 1% SDS(10min), IF-staining. Data obtained from customer.



EB06991 (0.5 $\mu$ g/ml) 48h-staining of PFA-perfused cryosection of Mouse Olfactory bulb. Antigen retrieval with citrate buffer pH 6 at 95C for 10min, IF-staining for NCAM2 (red), VGLUT2 (blue) and NQO1 (green), data obtained from customer.



EB06991 (3.75µg/ml) staining of paraffin embedded Human Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.