

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB11839 - Goat Anti-NCU-G1 (mouse aa243-256) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** 0610031J06Rik, AB027141, lysosomal protein NCU-G1, NCU-G1, RIKEN cDNA 0610031J06 gene

**Official Symbol:** Glmp

**Accession Number(s):** NP\_064387.1

**Non-Human GeneID(s):** 56700 (mouse), 295231 (rat)

**Important Comments:** The immunizing peptide represents the luminal part of the protein

### Immunogen

Peptide with sequence PDCPSVNERNSIDD, from the internal region of the protein sequence according to NP\_064387.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 75kDa band observed in Mouse and Rat Kidney lysates (calculated MW of 43.8kDa according to Mouse NP\_064387.1 and Rat NP\_001004226.1). The observed molecular weight corresponds to the endogenous wild-type glycoprotein, as shown in earlier findings in literature with different antibodies (Schieweck O et al, PMID:19489740). Recommended concentration: 0.3-1µg/ml.

### Species Reactivity

**Tested:** Mouse, Rat

**Expected from sequence similarity:** Mouse



EB11839 (0.5 $\mu$ g/ml) staining of Mouse (A) and Rat (B) Kidney lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.