

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB09788 - Goat Anti-NDUFS6 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: NADH dehydrogenase (ubiquinone) Fe-S protein 6, 13kDa

(NADH-coenzyme Q reductase), NDUFS6

Official Symbol: NDUFS6

Accession Number(s): NP_004544.1

Human GenelD(s): 4726

Non-Human GenelD(s): 407785 (mouse), 29478 (rat)

Immunogen

Peptide with sequence C-RIRFVGRQKEVNEN, from the internal region of the protein sequence according to NP_004544.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 15kDa band observed in Human Heart lysates and 14kDa in Mouse and Rat Heart lysates (calculated MW of 13.7kDa according to NP_004544.1). Recommended concentration: 0.01-0.03µg/ml. Primary incubation was 1 hour. **IHC:** Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.

Additional validation: This antibody has been successfully used in the following paper: Sikorski et al. (2018) PMID: 30377371.

Species Reactivity

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

Specific Reference


This antibody has been successfully used in the following paper:

Krzysztof Sikorski, Adi Mehta, Marit Inngjerdingen, Flourina Thakor, Simon Kling, Tomas Kalina, Tuula A. Nyman, Maria Ekman Stensland, Wei Zhou, Gustavo A. De Souza, Lars Holden, Jan Stuchly, Markus Templin and Fridtjof Lund-Johansen


A high-throughput pipeline for validation of antibodies


Nat Methods. 2018 Nov;15(11):909-912

PMID: 30377371



EB09788 (0.01µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.





EB09788 (5μg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.