

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB10194 - Goat Anti-NDUFS8 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** CI23KD, complex I-23kD, NADH dehydrogenase (ubiquinone) Fe-S protein 8, 23kDa (NADH-coenzyme Q reductase), NADH dehydrogenase ubiquinone Fe-S

8, NADH-ubiquinone oxidoreductase 23 kDa subunit, TYKY, NDUFS8

Official Symbol: NDUFS8

Accession Number(s): NP\_002487.1

Human GeneID(s): 4728

Non-Human GenelD(s): 225887 (mouse), 293652 (rat)

### **Immunogen**

Peptide with sequence C-AEPRADGSRR, from the Internal region of the protein sequence according to NP\_002487.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:2000.

**Western blot:** Approx 25kDa band observed in Human and Mouse Heart lysates (calculated MW of 23.7kDa according to NP\_002487.1). Recommended concentration: 0.1-0.3µg/ml. An additional fainter band of 90kDa was consistently observed, however this band was not blocked by the immunizing peptide and it is therefore a non-specific signal. We call for caution when used for other assays than Western blot.

# **Species Reactivity**

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB10194 (0.1 $\mu$ g/ml) staining of Mouse Heart lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.