

### **UK Office**

#### **Everest Biotech Ltd**

**Cherwell Innovation Centre** 

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

### **US Office**

### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB10193 - Goat Anti-NDUFS8 Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** CI23KD, complex I-23kD, NADH dehydrogenase (ubiquinone) Fe-S protein 8, 23kDa (NADH-coenzyme Q reductase), NADH dehydrogenase ubiquinone Fe-S

8, NADH-ubiquinone oxidoreductase 23 kDa subunit, TYKY, NDUFS8

Official Symbol: NDUFS8

Accession Number(s): NP\_002487.1

Human GeneID(s): 4728

Non-Human GenelD(s): 225887 (mouse), 293652 (rat)

### **Immunogen**

Peptide with sequence C-EAEIAANIQAD, from the internal region of the protein sequence according to NP\_002487.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:4000.

Western blot: Approx 25kDa band observed in Mouse and Rat Heart lysates (calculated

MW of 24.0kDa according to Mouse NP\_659119.2 and Rat NP\_001099792.1).

Recommended concentration: 0.1-0.3µg/ml.

### **Species Reactivity**

Tested: Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB10193 (0.1 $\mu$ g/ml) staining of Mouse Heart lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.