

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 **United States** 

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09686-B - Goat Anti-Neurogranin precursor, **Biotinylated Antibody**

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: NRGN, neurogranin, RC3, hng, calmodulin-binding protein, neurogranin (protein kinase C substrate, RC3), ng, protein kinase C substrate

Official Symbol: NRGN

Accession Number(s): NP\_006167.1

Human GenelD(s): 4900

Important Comments: Reported variants represent identical protein: NP\_006167.1,

NP\_001119653.1

### **Immunogen**

Peptide with sequence CKIKSGERGRKG., from the internal region of the protein sequence according to NP\_006167.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 16kDa band observed in Human Brain (Frontal Cortex) lysate (calculated MW of 7.6kDa according to NP\_006167.1). The observed molecular weights correspond to earlier findings in literature with different antibodies (Li et al, J Biol Chem. 1999 Jan 15;274(3):1294-300; PMID: 9880498). See non-biotinylated parental product's

datasheet for further QC data. Recommended concentration: 0.1-0.3µg/ml.

#### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Dog, Cow



Biotinylated EB09686 (0.1µg/ml) staining of Human Brain (Frontal Cortex) lysate (35µg protein in RIPA buffer).

Primary incubation was 1 hour. Detected by chemiluminescence.