

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06930 - Goat Anti-Neuroligin 4 / NLGN4X / NLGN4Y Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: NLGN4X, neuroligin 4, X-linked, HGNC:14287, AUTSX2, HLNX, HNLX, KIAA1260, MGC22376, NLGN, NLGN4, X-linked neuroligin 4, neuroligin X, NLGN4Y, neuroligin 4, Y-linked, KIAA0951, ASPGX2, OTTHUMP00000022864,

OTTHUMP00000034787, neuroligin 4, Y linked, neuroligin Y

Official Symbol: NLGN4X; NLGN4Y

Accession Number(s): NP_065793.1; NP_851849.1; NP_055708.3

Human GeneID(s): 57502, 22829

Important Comments: This antibody is expected to recognise both X-linked reported products (NP_065793.1 and NP_851849.1), which are identical, and the Y-linked reported product (NP_055708.3).

Immunogen

Peptide with sequence C-KHSKDPHKTGPED, from the internal region of the protein sequence according to NP_065793.1; NP_851849.1; NP_055708.3.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Preliminary experiments gave a faint approx 90kDa band and a strong approx 70kDa band in Human Brain lysates after 1μg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the 70kDa band we observe given the calculated size of 91.9kDa according to NP_065793.1, NP_851849.1 and .NP_055708.2 The 70kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

Species Reactivity

Tested:

Expected from sequence similarity: Human