

## **International Office**

Everest Biotech Ltd Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service: <u>customerservice@vectorlabs.com</u> Technical Service: <u>technical@vectorlabs.com</u>

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06769 - Goat Anti-Neuropeptide Y receptor Y5 Antibody



Size: 100µg specific antibody in 200µl

## **Target Protein**

Principal Names: neuropeptide Y receptor Y5, HGNC:7958, NPY5R, NPYR5, NPY5-R, NPY-Y5 receptor, NPYY5-R, Y5 receptor Official Symbol: NPY5R Accession Number(s): NP\_006165.1 Human GeneID(s): <u>4889</u> Non-Human GeneID(s): 18168 (mouse), 25340 (rat)

## Immunogen

Peptide with sequence C-GLSNKENRLEEN, from the internal region of the protein sequence according to NP\_006165.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 50kDa band observed in Human Brain, Breast and Duodenum lysates (calculated MW of 50.7kDa according to NP\_006165.1). Recommended concentration: 0.3-1µg/ml. Primary incubation was 1 hour. This antibody has been successfully used in WB on CHO-K1 cells: PMID: 30503694. This antibody has been successfully used in WB on Human: PMID: 28413811 and 20332211. **IHC:** In paraffin embedded Human Intestine shows staining of ganglion cells. Recommended concentration, 4-6µg/ml.

**Immunoprecipitation:** Customer feedback showed specific results when this product was used in IP. This antibody has been successfully used in IP: PMID: 30503694.

## **Species Reactivity**

Tested: Human, Hamster Expected from sequence similarity: Human, Mouse, Rat, Dog

## **Specific References**

This antibody has been successfully used in the following paper: Congyi Lu et al. Hypoxia-activated neuropeptide Y/Y5 receptor/RhoA pathway triggers chromosomal instability and bone metastasis in Ewing sarcoma. Nat Commun. 2022 Apr 28;13(1):2323. PMID: 35484119

#### This antibody has been successfully used in WB on Human:

Nouran Abualsaud, Lindsay Caprio, Susana Galli, Ewa Krawczyk, Lamia Alamri, Shiya Zhu, G Ian Kitlinska

Neuropeptide Y/Y5 Receptor Pathway Stimulates Neuroblastoma Cell Motility Through

RhoA Activation Front Cell Dev Biol. 2021 Feb 17;8:627090. PMID: 33681186

#### This antibody has been successfully used in WB and IP on CHO-K1 cells:

Magdalena Czarneckaa, Congyi Lua,b, Jennifer Ponsa, Induja Maheswarana, Pawel Ciborowskic, Lihua Zhangd, Amrita Cheemad, Joanna Kitlinskaa Neuropeptide Y receptor interactions regulate its mitogenic activity Neuropeptides. 2019 Feb;73:11-24. PMID: 30503694

#### This antibody has been successfully used in WB on Human:

Czarnecka M, Kitlinska J. Cell Surface Protein Detection to Assess Receptor Internalization. Bio Protoc. 2016 Oct 20;6(20) PMID: 28413811

#### This antibody has been successfully used in WB on Human:

Sheriff S, Ali M, Yahya A, Haider KH, Balasubramaniam A, Amlal H. Neuropeptide Y Y5 receptor promotes cell growth through extracellular signal-regulated kinase signaling and cyclic AMP inhibition in a human breast cancer cell line. Mol Cancer Res. 2010 Apr;8(4):604-14. PMID: 20332211



EB06769 (0.3µg/ml) staining of Human Brain lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB06769 (3.8µg/ml) staining of paraffin embedded Human Intestine. Steamed antigen retrieval with citrate buffer Ph 6, AP-staining.