

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09503 - Goat Anti-NFATC2 / NFAT1 Antibody

Size: 100µg specific antibody in 200µl



#### **Target Protein**

**Principal Names:** NFATC2, nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 2, KIAA0611, NFAT1, NFATP, NFAT pre-existing subunit, NFAT transcription complex, preexisting component, OTTHUMP00000031291, T cell transcription factor NFAT1, nuclear factor of activated T-cells, cytoplasmic 2, nuclear factor of activated T-cells, preexisting component, preexisting nuclear factor of activated T-cells 2

Official Symbol: NFATC2

Accession Number(s): NP\_036472.2; NP\_775114.1; NP\_001129493.1

Human GeneID(s): 4773

Non-Human GenelD(s): 140488 (mouse), 312320 (rat)

Important Comments: This antibody is expected to recognize all three reported isoforms

(NP\_036472.2; NP\_775114.1; NP\_001129493.1).

### **Immunogen**

Peptide with sequence C-QQFRTGLSSPDARYQ, from the internal region of the protein sequence according to NP\_036472.2; NP\_775114.1; NP\_001129493.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** Approx 110kDa band observed in Human lysates of cell line MOLT4 (calculated MW of 100kDa according to NP\_775114.1). Recommended concentration: 0.3-1µg/ml.

IHC: In paraffin embedded Human Spleen shows nuclear membrane and weaker cytoplasm staining in select splenocytes. Recommended concentration, 3-6µg/ml.


### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB09503 (0.3µg/ml) staining of MOLT4 lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB09503 (3.8µg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.