

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB05227 - Goat Anti-NFIL3 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: NFIL3, nuclear factor, interleukin 3 regulated, E4BP4, IL3BP1, NFIL3A, NF-IL3A, interleukin-3 promoter transcriptional activator, OTTHUMP00000021633

Official Symbol: NFIL3

Accession Number(s): NP_005375.2

Human GeneID(s): 4783

Non-Human GenelD(s): 18030 (mouse), 114519 (rat)

Immunogen

Peptide with sequence C-RLIATQPISASDSG, from the C Terminus of the protein sequence according to NP_005375.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: Approx. 60kDa band observed in nuclear lysates of cell line Jurkat and HeLa (calculated MW of 51.5kDa according to NP_005375.2). This molecular weight is routinely observed by other sources. An additional band of 30kDa was observed in lysates of cell line HeLa, which was completely blocked with the immunizing peptide.

Recommended concentration: 0.5-2µg/ml. Primary incubation 1 hour at room temperature.

Flow Cytometry: Flow cytometric analysis of Human peripheral blood mononuclear cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat



EB05227 staining (1µg/ml) of Jurkat nuclear cell lysate (RIPA buffer, 20µg total protein per lane). Detected using chemiluminescence.



EB05227 Flow cytometric analysis of paraformaldehyde fixed Human peripheral blood mononuclear cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.