

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07294 - Goat Anti-NMDAR2B / GRIN2B Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: GRIN2B, NMDAR2B, glutamate receptor, ionotropic, N-methyl D-aspartate 2B, HGNC:4586, NR2B, hNR3, N-methyl-D-aspartate receptor subunit 2B

Official Symbol: GRIN2B

Accession Number(s): NP_000825.1

Human GenelD(s): 2904

Non-Human GenelD(s): 14812 (mouse), 24410 (rat)

Immunogen

Peptide with sequence C-KDAHEKDDFHHLS, from the internal region of the protein sequence according to NP_000825.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 170kDa band observed in Rat Brain lysates (calculated MW of 166kDa according to NP_000825.1). Recommended concentration: 2-4µg/ml. Primary incubation was 1 hour.

Immunofluorescence: Strong expression of the protein seen in the cytoskeleton/cytoplasm of HeLa cells. Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Human, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 15kDa

EB07294 (2μg/ml) staining of Rat Brain lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB07294 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoskeleton and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).