



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11023 - Goat Anti-NMNAT3 (aa149-162) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: NMNAT3, nicotinamide nucleotide adenylyltransferase 3, PNAT3, NMN adenylyltransferase 3, NaMN adenylyltransferase 3, nicotinamide mononucleotide adenylyltransferase 3, nicotinate-nucleotide adenylyltransferase 3, pyridine nucleotide adenylyltransferase 3, NaMN adenylyltransferase 3, nicotinamide mononucleotide adenylyltransferase 3, nicotinamide nucleotide adenylyltransferase 3, nicotinate-nucleotide adenylyltransferase 3, NMN adenylyltransferase 3, PNAT3, pyridine nucleotide adenylyltransferase 3, NMNAT3

Official Symbol: NMNAT3

Accession Number(s): NP_835471.1; NP_001186976.1; NP_001307439.1;
NP_001307440.1; NP_001307442.1

Human GeneID(s): [349565](#)

Important Comments: This antibody is expected to recognize all reported isoforms

Immunogen

Peptide with sequence C-QHNIHLAKEPVQNE, from the internal region of the protein sequence according to NP_835471.1; NP_001186976.1; NP_001307439.1; NP_001307440.1; NP_001307442.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Preliminary testing showed a band at approx. 26kDa in Caco-2, A431 and A549 cell lysate at a concentration of 2µg/ml (calculated Mwt. of 24.1kDa according to NP_835471.1). Primary incubation 1 hour at room temperature.


Immunofluorescence: Strong expression of the protein seen in HEK293 cells.
Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of HEK293 cells. Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog



EB11023 Immunofluorescence analysis of paraformaldehyde fixed HEK293 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic and some nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB11023 Flow cytometric analysis of paraformaldehyde fixed HEK293 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.