



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07246 - Goat Anti-NONO / p54NRB Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: NONO, P54NRB, non-POU domain containing, octamer-binding, HGNC:7871, NMT55, NRB54, P54, Nuclear RNA-binding protein, 54-kD, non-POU-domain-containing, octamer-binding, non-Pou domain-containing octamer (ATGCAAAT) binding protein

Official Symbol: NONO

Accession Number(s): NP_031389.3

Human GeneID(s): [4841](#)

Non-Human GeneID(s): 53610 (mouse), 317259 (rat)

Immunogen

Peptide with sequence C-NRAAPGAEFAPNK, from the C Terminus of the protein sequence according to NP_031389.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.


Western blot: Approx 60kDa band observed in Mouse Brain lysates (calculated MW of 54.2kDa according to Human NP_031389.3 and 54.5kDa according to Mouse NP_075633.2). Recommended concentration: 0.1-0.3µg/ml.

IHC: Paraffin embedded Human Spleen and Testis. Recommended concentration: 3.75µg/ml.


Species Reactivity

Tested: Human, Mouse


Expected from sequence similarity: Human, Mouse, Rat, Dog



EB07246 (0.1µg/ml) staining of Mouse Brain lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB07246 (3.75µg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB07246 (3.75µg/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.