



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05259-T - Goat Anti-NOS1 Antibody - Trial

Size: 20µg specific antibody in 40µl



Target Protein

Principal Names: NOS1, nitric oxide synthase 1 (neuronal), NOS, neuronal nitric oxide synthase, PnNOS, penile neuronal nitric oxide synthase, penile neuronal NOS, IHPS1, nNOS, nitric oxide synthase 1, neuronal

Official Symbol: NOS1

Accession Number(s): NP_000611.1; NP_001191147.1; NP_001191142.1

Human GeneID(s): [4842](#)

Immunogen

Peptide with sequence C-ESKKDTDEVFSS, from the C Terminus of the protein sequence according to NP_000611.1; NP_001191147.1; NP_001191142.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx. 160kDa band observed in Mouse Brain lysates (calculated MW of 160.5kDa according to NP_032738.1). Recommended concentration: 1-2µg/ml. Primary incubation 1 hour at room temperature.

IHC: Cryosections of Human Hypothalamus. Recommended dilution range: 1:10000 - 1:40000.

Immunofluorescence: Strong expression of the protein seen in the nuclei of HeLa and U2OS cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of HeLa cells. Recommended concentration: 10ug/ml.

Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

Specific Reference


This antibody (previous batch) has been successfully used in IF on Mouse:

Fried HU, Kaupp UB, Müller F.


Hyperpolarization-activated and cyclic nucleotide-gated channels are differentially expressed in juxtglomerular cells in the olfactory bulb of mice.

Cell Tissue Res. 2010 Mar;339(3):463-79.


PMID: 20140458




EB05259 (1 μ g/ml) staining of Mouse Brain lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.




EB05259 Immunostaining of 30 μ m thick cryosections of PFA-perfused Human Hypothalamus, antigen retrieval with citrate buffer Ph 6 at 80C for 30 min, HRP-staining with Ni-DAB after Biotin-SP-antigoat amplification. Data obtained by Drs. Szabolcs Takács and Erik Hrabovszky, Inst, Exp, Med, Budapest, Hungary.



EB05259 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB05259 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB05259 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.