

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB10571 - Goat Anti-NPAS4 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** NPAS4, neuronal PAS domain protein 4, Le-PAS, NXF, PASD10, bHLHe79, HLH-PAS transcription factor NXF, PAS domain-containing protein 10, class E

basic helix-loop-helix protein 79, neuronal PAS4

Official Symbol: NPAS4

Accession Number(s): NP\_849195.2

Human GeneID(s): 266743

Non-Human GenelD(s): 225872 (mouse), 266734 (rat)

## **Immunogen**

Peptide with sequence CRFNTSKSLRRQS, from the internal region of the protein sequence according to NP\_849195.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:4000.

**Western blot:** Approx 90kDa band observed in Human Brain (Frontal Cortex) lysates and in Mouse and Rat Brain lysates (calculated MW of 87.1kDa according to NP\_849195.2).

Recommended concentration: 0.3-1µg/ml.


### **Species Reactivity**

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB10571 (0.3 $\mu$ g/ml) staining of Human Frontal Cortex lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB10571 (1μg/ml) staining of Mouse (A) and Rat (B) Brain lysates (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.