

International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09947 - Goat Anti-NPC2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: EDDM1, epididymal protein 1, epididymal secretory protein, epididymal secretory protein E1, HE1, MGC1333, Niemann-Pick disease, type C2, NP-C2, tissue-specific secretory protein, NPC2

Official Symbol: NPC2

Accession Number(s): NP_006423.1

Human GeneID(s): [10577](#)

Immunogen

Peptide with sequence QAEPVQFKDCGSVD, from the internal region of the protein sequence according to NP_006423.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: Experiments gave bands at approx 20kDa and 23kDa in Human Testis lysates. These bands correspond to earlier findings in literature with different antibodies (Légaré et al, Mol Hum Reprod. 2006 Jul;12(7):461-8; PMID: 16772431. This protein has a calculated MW of 16.6kDa according to NP_006423.1. Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB09947 (0.5µg/ml) staining of Human Testis lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.