



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11190 - Goat Anti-NPR3 (aa520-533) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ANP-C, ANPRC, ANPR-C, atrial natriuretic peptide clearance receptor, atrial natriuretic peptide receptor 3, atrial natriuretic peptide receptor type C, GUCY2B, natriuretic peptide receptor C/guanylate cyclase C (atrionatriuretic peptide receptor C), NPRC, NPR-C, OTTHUMP00000197048, OTTHUMP00000220980, NPR3

Official Symbol: NPR3

Accession Number(s): NP_000899.1

Human GeneID(s): [4883](#)

Non-Human GeneID(s): 18162 (mouse), 25339 (rat)

Important Comments: The immunizing peptide represents the cytoplasmic part of the protein.

Immunogen

Peptide with sequence C-HRELREDSIRSHFS, from the C Terminus of the protein sequence according to NP_000899.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 52kDa band observed in Human Placenta lysates (calculated MW of 59.8kDa according to NP_000899.1). Recommended concentration: 1-3µg/ml.


IHC: In paraffin embedded Human Kidney shows staining of brush borders in select PCT. Recommended concentration: 2-4µg/ml. Paraffin embedded Human Prostate.

Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow




EB11190 (1 μ g/ml) staining of Human Placenta lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB11190 (2 μ g/ml) staining of paraffin embedded Human Kidney. Microwaved antigen retrieval with citrate buffer pH 6, HRP-staining.



EB11190 Negative Control showing staining of paraffin embedded Human Kidney, with no primary antibody.



EB11190 (10µg/ml) staining of paraffin embedded Human Prostate. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.