

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07390 - Goat Anti-NTRK2 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** NTRK2, neurotrophic tyrosine kinase, receptor, type 2, GP145-TrkB, TRKB, BDNF/NT-3 growth factors receptor, tyrosine kinase receptor B, BDNF receptor

Official Symbol: NTRK2

Accession Number(s): NP\_006171.2; NP\_001007098.1; NP\_001018074.1;

NP\_001018075.1; NP\_001018076.1

Human GeneID(s): 4915

**Important Comments:** This antibody is expected to recognise isoforms a (NP\_006171.2), b ((NP\_001007098.1), c (NP\_001018074.1), d (NP\_001018075.1), e ( NP\_001018076.1).

### **Immunogen**

Peptide with sequence C-KTLQEAKSSPDTQ, from the internal region of the protein sequence according to NP\_006171.2; NP\_001007098.1; NP\_001018074.1; NP\_001018075.1; NP\_001018076.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:64000.


**Western blot:** Approx 100kDa band observed in Human Brain (Hippocampus, Cerebral Cortex and Cerebellum) lysates (calculated MW of 93.8kDa according to NP\_006171.2).

Recommended concentration: 0.05-0.2µg/ml.

#### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB07390 (0.05µg/ml) staining of Human Hippocampus (A), Cerebral Cortex (B) and Cerebellum (C) lysates (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.