

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09516 - Goat Anti-NUDC Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** NUDC, nuclear distribution gene C homolog (A. nidulans), HNUDC, MNUDC, NPD011, OTTHUMP0000004405, nuclear distribution gene C homolog

Official Symbol: NUDC

Accession Number(s): NP\_006591.1

Human GenelD(s): 10726

Non-Human GenelD(s): 18221 (mouse), 29648 (rat)

### **Immunogen**

Peptide with sequence C-PNLGNGADLPNYR, from the internal region of the protein sequence according to NP\_006591.1.

Please note the <u>peptide</u> is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.


**Western blot:** Approx 40kDa band observed in lysates of cell lines HeLa. HepG2 and K562, and approx. 38kDa in lysates of cell line KNRK (calculated MW of 38.2kDa according to Human NP\_006591.1) and 38.4kDa according to Rat NP\_058967.1). Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature.

**Immunofluorescence:** Strong expression of the protein seen in the cytoplasm of A431 cells. Recommended concentration: 10µg/ml.


#### **Species Reactivity**

Tested: Human, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB09516 (0.5μg/ml) staining of HeLa (A), HepG2 (B), K562 (C) and KNRK (D) cell lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB09516 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa

Fluor 488 secondary antibody (2ug/ml).