



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB10160 - Goat Anti-NUP107 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: nucleoporin 107kDa, NUP84, NUP107

Official Symbol: NUP107

Accession Number(s): NP_065134.1

Human GeneID(s): [57122](#)

Non-Human GeneID(s): 103468 (mouse), 116555 (rat)

Immunogen

Peptide with sequence C-PLPAEDDNAIREH, from the internal region of the protein sequence according to NP_065134.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.


Western blot: Approx 100kDa band observed in Human Peripheral Blood

Mononucleocytes lysates (calculated MW of 106kDa according to Human NP_065134.1) and 120kDa in nuclear lysates of NIH3T3 (calculated MW of 107kDa according to Mouse NP_598771.1). Recommended concentration: 1-3µg/ml.


Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB10160 (1 μ g/ml) staining of Peripheral Blood Lymphocytes lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB10160 (2 μ g/ml) staining of nuclear lysate of NIH3T3 (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.