

#### **UK Office**

### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$ 

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB12375 - Goat Anti-OAS3 Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** OAS3, 2'-5'-oligoadenylate synthetase 3, 100kDa, p100, (2-5')oligo(A) synthase 3, (2-5')oligo(A) synthetase 3, 2'-5'-oligoadenylate synthase 3,

2'-5'oligoadenylate synthetase p100, 2-5A synthase 3, 2-5A synthetase 3, p100 OAS,

p100OAS

Official Symbol: OAS3

Accession Number(s): NP\_006178.2

Human GeneID(s): 4940

## **Immunogen**

Peptide with sequence C-HPIQLDPNQKTPENS, from the internal region of the protein sequence according to NP\_006178.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** Approx 110kDa band observed in lysates of cell line A431 (calculated MW of 121kDa according to NP\_006178.2). Recommended concentration: 1-3μg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa

> 25kDa 20kDa

15kDa

EB12375 ( $2\mu g/ml$ ) staining of A431 lysate ( $35\mu g$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.