



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05067 - Goat Anti-ORC4L Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ORC4L, origin recognition complex, subunit 4-like (yeast), Orc4p, HSORC4, origin recognition complex, subunit 4 (yeast homolog)-like, Origin recognition complex, subunit 4, S. cerevisiae, homolog-like, ORC4, ORC4P, origin recognition complex subunit 4, origin recognition complex, subunit 4 homolog

Official Symbol: ORC4

Accession Number(s): NP_002543.2; NP_001177810.1; NP_001177811.1

Human GeneID(s): [5000](#)

Non-Human GeneID(s): 26428 (mouse), 295596 (rat)

Important Comments: This antibody is expected to recognize all reported isoforms (NP_002543.2; NP_001177810.1; NP_001177811.1). Reported variants represent identical protein: NP_002543.2, NP_001177808.1, NP_859526.1, NP_859525.1

Immunogen

Peptide with sequence C-DVRQWATSSLSWL, from the C Terminus of the protein sequence according to NP_002543.2; NP_001177810.1; NP_001177811.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Approx 48kDa band observed in nuclear lysates of cell line Jurkat (calculated MW of 50.4kDa according to Human NP_002543.2 Recommended concentration: 0.3-1µg/ml. Primary incubation was 1 hour. Preliminary testing was unsuccessful on cell line NIH-3T3 and Mouse Heart for this particular batch.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB05067 (0.3 μ g/ml) staining of Jurkat nuclear lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.