

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05067-B - Goat Anti-ORC4L, Biotinylated Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: HSORC4, ORC4, origin recognition complex subunit 4, ORC4L, ORC4P, origin recognition complex, subunit 4 homolog, origin recognition complex, subunit 4 (yeast homolog)-like, Origin recognition complex, subunit 4, S. cerevisiae, homolog-like, origin recognition complex, subunit 4-like (yeast)

Official Symbol: ORC4

Accession Number(s): NP_002543.2; NP_001177810.1; NP_001177811.1

Human GeneID(s): [5000](#)

Non-Human GeneID(s): 26428 (mouse), 295596 (rat)

Important Comments: This antibody is expected to recognize all reported isoforms (NP_002543.2; NP_001177810.1; NP_001177811.1). Reported variants represent identical protein: XP_011509557.1, NP_002543.2, NP_001177808.1, NP_859526.1, NP_859525.1 Reported variants represent identical protein: XP_006712619.1, NP_001177810.1

Immunogen

Peptide with sequence C-DVRQWATSSLSWL., from the C Terminus of the protein sequence according to NP_002543.2; NP_001177810.1; NP_001177811.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.


Western blot: Approx 48kDa band observed in Jurkat nuclear lysates (calculated MW of 50.4kDa according to NP_002543.2). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.1-0.3µg/ml.

Chromatin immunoprecipitation: reported by a customer to work very well in ChIP.

Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



Biotinylated EB05067 (0.1µg/ml) staining of Jurkat nuclear lysate (35µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.