



## UK Office

**Everest Biotech Ltd**  
Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

**Everest Biotech c/o Abcore**  
405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09023 - Goat Anti-OXT Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** OXT, oxytocin, prepro- (neurophysin I), MGC126890, MGC126892, OT, OT-NPI, oxytocin-neurophysin I, oxytocin-neurophysin I, preproprotein

**Official Symbol:** OXT

**Accession Number(s):** NP\_000906.1

**Human GeneID(s):** [5020](#)

### Immunogen

Peptide with sequence HADPACDAEATFSQR, from the C Terminus of the protein sequence according to NP\_000906.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:4000.


**Western blot:** Preliminary experiments in Human Brain (Cerebellum, Amygdala) lysates gave no specific signal but low background (at antibody concentration up to 1µg/ml). We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates?

**IHC:** Frozen section of Human Hypothalamus shows staining of magnocellular oxytocin neurons in the supraoptic nucleus. Recommended concentration, 0.05-0.15µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB09023 (0.05 $\mu$ g/ml) staining of PFA-perfused cryosection of Human Hypothalamus. Antigen retrieval with citrate buffer pH 6 at 80C for 30min, HRP-staining with Ni-DAB after Biotin-SP-antigoat amplification. Data obtained by Prof. Erik Hrabovszky, Inst, Exp, Med., Budapest, Hungary.