



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09509 - Goat Anti-P2RX7 / P2X7 receptor Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** P2RX7, purinergic receptor P2X, ligand-gated ion channel, 7, MGC20089, P2X7, ATP receptor, P2X purinoceptor 7, P2X7 receptor, P2Z receptor, purinergic receptor P2X7

**Official Symbol:** P2RX7

**Accession Number(s):** NP\_002553.3

**Human GeneID(s):** [5027](#)

**Non-Human GeneID(s):** 18439 (mouse), 29665 (rat)

### Immunogen

Peptide with sequence YETNKVTRIQSMNY-C, from the N Terminus of the protein sequence according to NP\_002553.3.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx 37kDa and 70kDa band observed in Human Brain (Frontal Cortex) lysates (calculated MW of 68.6kDa according to NP\_002553.2). Recommended concentration: 1-3µg/ml. Primary incubation was an 1 hour. An additional band of unknown identity was also consistently observed at 37kDa. This band was successfully blocked by incubation with the immunizing peptide.

**IHC:** In paraffin embedded Human Brain Cortex shows membranous staining of cell bodies and processes. Recommended concentration: 3-6µg/ml. Paraffin embedded Human Kidney. Recommended concentration: 3.75µg/ml.

**Immunofluorescence:** Strong expression of the protein seen in the cytoplasm and plasma membranes of HeLa cells. Recommended concentration: 10µg/ml. This antibody has been successfully used in IF on Mouse: Thaler R et al. (2015) PMID: 25491310.

**Flow Cytometry:** Flow cytometric analysis of HeLa cells. Recommended concentration: 10ug/ml.

### Species Reactivity

**Tested:** Human, Mouse

**Expected from sequence similarity:** Human, Mouse, Rat, Dog

### Specific Reference

**This antibody has been successfully used in IF on Mouse:**


Thaler R, Sturmlechner I, Spitzer S, Riester SM, Rumpler M, Zwerina J, Klaushofer K, van Wijnen AJ, Varga F.

Acute-phase protein serum amyloid A3 is a novel paracrine coupling factor that controls bone homeostasis.


FASEB J. 2015 Apr;29(4):1344-59.  
PMID: 25491310




EB09509 (0.3µg/ml) staining of A431 (A) and HeLa (B) cell lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.




EB09509 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic and plasma membrane staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB09509 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



EB09509 (3.75 $\mu$ g/ml) staining of paraffin embedded Human Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB09509 (3.75µg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.