



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB08374 - Goat Anti-PAM / PAL Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** PAM, PAL, peptidylglycine alpha-amidating monooxygenase, PHM, pancreatic peptidylglycine alpha-amidating monooxygenase, peptidyl alpha-amidating enzyme, peptidyl-alpha-hydroxyglycine alpha-amidating lyase, peptidylglycine 2-hydroxylase, peptidylglycine alpha-hydroxylating monooxygenase

**Official Symbol:** PAM

**Accession Number(s):** NP\_000910.2; NP\_620121.1; NP\_620176.1; NP\_620177.1

**Human GeneID(s):** [5066](#)

**Important Comments:** This antibody is expected to recognise all reported isoforms (NP\_000910.2; NP\_620121.1; NP\_620176.1 and NP\_620177.1).

### Immunogen

Peptide with sequence C-DQEKEDDGSESEE, from the C Terminus of the protein sequence according to NP\_000910.2; NP\_620121.1; NP\_620176.1; NP\_620177.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:4000.

**Western blot:** Approx 100kDa band observed in Human Kidney lysates (calculated MW of 101kDa according to preproprotein NP\_620121.1). Recommended concentration: 0.3-1µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB08374 (1µg/ml) staining of Human Kidney lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.  
Detected by chemiluminescence.