



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB13107 - Goat Anti-pan ADH (N terminus) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** pan ADH, ADH1B, ADH2, alcohol dehydrogenase IB (class I), beta polypeptide, ADH, beta subunit, aldehyde reductase, alcohol dehydrogenase 2, alcohol dehydrogenase 2 (class I), beta polypeptide, alcohol dehydrogenase 1B (class I), beta polypeptide

**Official Symbol:** ADH1A, B, C

**Accession Number(s):** NP\_000659.2; NP\_000658.1; NP\_000660.1

**Human GeneID(s):** [124](#) , [125](#)

**Important Comments:** This is an alternative product to EB06599. This antibody is expected to recognise the alpha (ADH1A, NP\_000658.1), the beta (ADH1B, NP\_000659.2) and gamma (ADH1C, NP\_000660.1) polypeptide variants of human alcohol dehydrogenase.

### Immunogen

Peptide with sequence STAGKVIKCKAC, from the N Terminus of the protein sequence according to NP\_000659.2; NP\_000658.1; NP\_000660.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:4000.

**Western blot:** Approx 38kDa band observed in Human Liver, Duodenum and Breast lysates and approx. 37kDa in Rat Liver lysates (calculated MW of 39.9kDa according to Human NP\_000659.2 and 39.6kDa Rat NP\_062159.3). Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature.


### Species Reactivity

**Tested:** Human, Rat

**Expected from sequence similarity:** Human, Mouse, Rat



EB13107 (0.3ug/ml) staining of Human Liver (A), (0.5ug/ml) Duodenum (B) and (1ug/ml) Breast (C) lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB13107 (0.5ug/ml) staining of Rat Liver lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.