



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB13107 - Goat Anti-pan ADH (N terminus) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: pan ADH, ADH1B, ADH2, alcohol dehydrogenase IB (class I), beta polypeptide, ADH, beta subunit, aldehyde reductase, alcohol dehydrogenase 2, alcohol dehydrogenase 2 (class I), beta polypeptide, alcohol dehydrogenase 1B (class I), beta polypeptide

Official Symbol: ADH1A, B, C

Accession Number(s): NP_000659.2; NP_000658.1; NP_000660.1

Human GeneID(s): [124](#) , [125](#)

Important Comments: This is an alternative product to EB06599. This antibody is expected to recognise the alpha (ADH1A, NP_000658.1), the beta (ADH1B, NP_000659.2) and gamma (ADH1C, NP_000660.1) polypeptide variants of human alcohol dehydrogenase.

Immunogen

Peptide with sequence STAGKVIKCKAC, from the N Terminus of the protein sequence according to NP_000659.2; NP_000658.1; NP_000660.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:4000.

Western blot: Approx 38kDa band observed in Human Liver, Duodenum and Breast lysates and approx. 37kDa in Rat Liver lysates (calculated MW of 39.9kDa according to Human NP_000659.2 and 39.6kDa Rat NP_062159.3). Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature.


Species Reactivity

Tested: Human, Rat

Expected from sequence similarity: Human, Mouse, Rat



EB13107 (0.3ug/ml) staining of Human Liver (A), (0.5ug/ml) Duodenum (B) and (1ug/ml) Breast (C) lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB13107 (0.5ug/ml) staining of Rat Liver lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.