



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06236 - Goat Anti-PAPIN / PDZK3 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PDZD2, PDZK3, PAPIN, KIAA0300, PDZ domain containing 3, PDZ domain containing 2, likely ortholog of rat plakophilin-related armadillo repeat protein-interacting PDZ protein, PDZ domain containing 2, AIPC, PIN1, activated in prostate cancer protein, plakophilin-related armadillo repeat protein-interacting PDZ protein-like

Official Symbol: PDZD2

Accession Number(s): NP_835260.2

Human GeneID(s): [23037](#)

Immunogen

Peptide with sequence PITQDNAVLHLPLC, from the N Terminus of the protein sequence according to NP_835260.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


IHC: Paraffin embedded Human Tonsil. Recommended concentration: 6µg/ml.

Immunofluorescence: Expression of the protein seen in the plasma membrane and cytoplasm of U2OS cells. Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human



EB06236 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing plasma membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06236 (6µg/ml) staining of paraffin embedded Human Tonsil. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB06236 Negative Control showing staining of paraffin embedded Human Tonsil, with no primary antibody.