



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

### Everest Biotech c/o Abcore

405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB07110 - Goat Anti-PARP12 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** PARP12, poly (ADP-ribose) polymerase family, member 12, HGNC:21919, FLJ22693, PARP-12, ZC3H1, ZC3HDC1, zinc finger CCCH type domain containing 1, zinc finger CCCH-type domain containing 1

**Official Symbol:** PARP12

**Accession Number(s):** NP\_073587.1

**Human GeneID(s):** [64761](#)

### Immunogen

Peptide with sequence C-NAHDIKNKSSAP, from the internal region of the protein sequence according to NP\_073587.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:128000.

**Western blot:** Preliminary experiments gave an approx 110kDa band in Human Liver lysates and in lysates of hepatoblastoma cell line HEPG2 after 0.03µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 79.1kDa according to NP\_073587.1. The 110kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

**Immunofluorescence:** Anonymous customer has confirmed specificity by siRNA-mediated PARP2 knockdown.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Dog