



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05513 - Goat Anti-PAX8 (N Terminus) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PAX8, paired box gene 8, paired domain gene 8, paired box 8

Official Symbol: PAX8

Accession Number(s): NP_003457.1; NP_039245.1; NP_039246.1; NP_039247.1;
NP_054698.1

Human GeneID(s): [7849](#)

Non-Human GeneID(s): 18510 (mouse), 81819 (rat)

Important Comments: This antibody is expected to recognise all five human isoforms (as represented by NP_003457.1; NP_039245.1; NP_039246.1; NP_039247.1; NP_054698.1)

Immunogen

Peptide with sequence PHNSIRSGHGGLN-C, from the N Terminus of the protein sequence according to NP_003457.1; NP_039245.1; NP_039246.1; NP_039247.1; NP_054698.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.


Western blot: Approx 48kDa band observed in Human Liver lysates. An additional 35kDa band was also observed in some Liver and HEK293 cell lysates (calculated MW of 34.9kDa according to Human NP_039247.1 PAX8D and 48.2kDa according to Human NP_003457.1 PAXA). Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Kidney. Recommended concentration: 4-5µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human, Mouse, Rat



EB05513 (1 μ g/ml) staining of Human Liver (A) and HEK293 cell (B) lysate (35 μ g protein in RIPA buffer).
Detected by chemiluminescence.



EB05513 (4 μ g/ml) staining of paraffin embedded Human Kidney. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB05513 Negative Control showing staining of paraffin embedded Human Kidney, with no primary antibody.