

## **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06944 - Goat Anti-PCK2 / PEPCK-M Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** PCK2, PEPCK-M, phosphoenolpyruvate carboxykinase 2 (mitochondrial), HGNC:8725, PEPCK, PEPCK2, PEP carboxykinase, mitochondrial phosphoenolpyruvate carboxykinase 2, phosphoenolpyruvate carboxylase,

phosphopyruvate carboxylase

Official Symbol: PCK2

Accession Number(s): NP\_004554.3; NP\_001278485.1

Human GeneID(s): 5106

Important Comments: No cross-reactivity with PCK1.

### Immunogen

Peptide with sequence KPWKPGDKEPCAH, from the internal region of the protein sequence according to NP\_004554.3; NP\_001278485.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:4000.

**Western blot:** Approx 70kDa band observed in lysates of cell lines A431, HEK293 HepG2 and NIH3T3 (calculated MW of 70.7kDa according to Human NP\_004554.3 and 70.5kDa according to Mouse NP\_083270.2). Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.

**Immunofluorescence:** Strong expression of the protein seen in the cytoplasm of U2OS cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of MCF7 cells. Recommended concentration: 10ug/ml.

# **Species Reactivity**

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

#### **Specific References**

#### This antibody (previous batch) has been successfully used in WB and IF on Human:

Méndez-Lucas A, Hyrossová P, Novellasdemunt L, Viñals F, Perales JC.

Mitochondrial phosphoenolpyruvate carboxykinase (PEPCK-M) is

a pro-survival, endoplasmicreticulum (ER) stress response gene involved in tumor cell adaptation to J Biol Chem. 2014 Aug 8;289(32):22090-102

PMID: 24973213

#### This antibody (previous batch) has been successfully used in WB and IF on Mouse:


Méndez-Lucas A, Duarte JA, Sunny NE, Satapati S, He T, Fu X, Bermúdez J, Burgess SC, Perales JC.

PEPCK-M expression in mouse liver potentiates, not replaces, PEPCK-C mediated

gluconeogenesis.

J Hepatol. 2013 Mar 4. doi:pii: S0168-8278(13)00142-6.


PMID: 23466304




EB06944 (2μg/ml) staining of A431 (A), HEK293 (B) and (1ug/ml) HepG2 (C) cell lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



.EB06944 (2μg/ml) staining of NIH3T3 cell lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB06944 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06944 Flow cytometric analysis of paraformaldehyde fixed MCF7 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control:

Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.