

## **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB11651-B - Goat Anti-PCNA (aa111-122), Biotinylated Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** cyclin, DNA polymerase delta auxiliary protein, MGC8367, OTTHUMP00000030189, OTTHUMP00000030190, PCNA, proliferating cell nuclear antigen

Official Symbol: PCNA

Accession Number(s): NP\_002583.1

Human GeneID(s): 5111

Non-Human GenelD(s): 18538 (mouse), 25737 (rat)

Important Comments: Reported variants represent identical protein: NP\_872590.1,

NP\_002583.1

## **Immunogen**

Peptide with sequence C-EAPNQEKVSDYE., from the internal region of the protein sequence according to NP\_002583.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 30kDa band observed in lysates of cell line Jurkat (calculated MW of 28.8kDa according to Human NP\_002583.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Gilljam et al, , J Cell Biol. 2009 Sep 7;186(5):645-54.). Recommended concentration: 0.03-0.1μg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



Biotinylated EB11651 (0.05ug/ml) staining of Jurkat lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.