



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07514 - Goat Anti-PCSK9 (aa164-175) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PCSK9, proprotein convertase subtilisin/kexin type 9, FH3, HCHOLA3, NARC-1, NARC1, hypercholesterolemia, autosomal dominant 3, neural apoptosis regulated convertase 1

Official Symbol: PCSK9

Accession Number(s): NP_777596.2

Human GeneID(s): [255738](#)

Immunogen

Peptide with sequence C-PRYRADEYQPPD, from the internal region of the protein sequence according to NP_777596.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: In transfected HEK293 transiently expressing full-length Human PCSK9 (myc and DYKDDDDK tagged), a doublet of approx. 80-100kDa was observed. No bands were observed in mock-transfected HEK293 and the same bands were observed using anti-myc tag antibody. Recommended concentration, 0.1-0.5µg/ml.

Purchase the +ve control! The transfected lysate used to QC this product is available for sale. Please email sales@everestbiotech.com to request datasheet, pricing and delivery information.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



HEK293 lysate (10ug protein in RIPA buffer) overexpressing Human PCSK9 with C-terminal MYC tag probed with EB07514 (0.5ug/ml) in Lane A and probed with anti-MYC Tag (1/1000) in lane C. Mock-transfected HEK293 probed with EB07514 (1mg/ml) in Lane B. Primary incubations were for 1 hour. Detected by chemiluminescence.