



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06682 - Goat Anti-PCSK9 (C Terminus) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** PCSK9, FH3, NARC1, HCHOLA3, proprotein convertase subtilisin/kexin type 9, neural apoptosis regulated convertase 1, hypercholesterolemia, autosomal dominant 3, proprotein convertase subtilisin/kexin type 9, LDLQC1, NARC-1

**Official Symbol:** PCSK9

**Accession Number(s):** NP\_777596.2

**Human GeneID(s):** [255738](#)

### Immunogen

Peptide with sequence CRSRHQAQASQELQ, from the C Terminus of the protein sequence according to NP\_777596.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx 75kDa band observed in Human Adipose lysates (calculated MW of 74.3kDa according to NP\_777596.2). Recommended concentration: 0.3-1µg/ml.

Primary incubation was 1 hour. Constructs and lysates were as described in Sun et al Hum Mol Genet. 2005 May 1;14(9):1161-9, PMID: 15772090. A customer found positive results in WB on Human Liver.

**IHC:** Paraffin embedded Human Liver. Recommended concentration: 5-6µg/ml.


### Species Reactivity

**Tested:** Human


**Expected from sequence similarity:** Human



EB06682 (0.3 $\mu$ g/ml) staining of Human Adipose lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.



EB06682 (5 $\mu$ g/ml) staining of paraffin embedded Human Liver. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB06682 Negative Control showing staining of paraffin embedded Human Liver, with no primary antibody.