



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

### Everest Biotech c/o Abcore

405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06201 - Goat Anti-Pericentrin 1 / NUP85 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** NUP85, pericentrin 1, PCNT1, PCN, PCNT, frount, nucleoporin Nup75, nucleoporin 85kDa, FLJ12549, Nup75, nucleoporin 85

**Official Symbol:** NUP85

**Accession Number(s):** NP\_079120.1; NP\_001290205.1; NP\_001317401.1

**Human GeneID(s):** [79902](#)

**Non-Human GeneID(s):** 18541 (mouse)

**Important Comments:** This antibody is expected to recognise all reported isoforms.

### Immunogen

Peptide with sequence C-ARAIIREGSLEGS, from the C Terminus of the protein sequence according to NP\_079120.1; NP\_001290205.1; NP\_001317401.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 75kDa band observed in HeLa lysates (calculated MW of 75kDa according to NP\_079120.1). Recommended concentration: 1-3µg/ml. Primary incubation was 1 hour.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat

### Specific References

**This antibody has been successfully used in IHC on Rat:**

Toyo Ikebukuro et al.

Disulfiram Ophthalmic Solution Inhibited Macrophage Infiltration by Suppressing Macrophage Pseudopodia Formation in a Rat Corneal Alkali Burn Model

Int. J. Mol. Sci. 2023, 24(1), 735; <https://doi.org/10.3390/ijms24010735>

PMID: 0

**This antibody has been successfully used in WB and IP on Human:**

Toda E, Terashima Y, Sato T, Hirose K, Kanegasaki S, Matsushima K.

FROUNT is a common regulator of CCR2 and CCR5 signaling to control directional migration.

J Immunol. 2009 Nov 15;183(10):6387-94.

PMID: 19841162



EB06201 (1µg/ml) staining of Hela lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.