



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06764 - Goat Anti-PGC-1beta / PERC Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PPARGC1B, PERC, PGC1B, PGC-1(beta), peroxisome proliferative activated receptor, gamma, coactivator 1, beta, PGC-1-related estrogen receptor alpha coactivator, peroxisome proliferator-activated receptor gamma coactivator 1 beta

Official Symbol: PPARGC1B

Accession Number(s): NP_573570.2

Human GeneID(s): [133522](#)

Non-Human GeneID(s): 170826 (mouse), 291567 (rat)

Immunogen

Peptide with sequence AGNDCGALLDEELS, from the N Terminus of the protein sequence according to NP_573570.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:4000.

IHC: In paraffin embedded Human Skeletal Muscle shows nuclear staining in muscle fibre cells. Recommended concentration, 3-6µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB06764 (3.8µg/ml) staining of paraffin embedded Human Skeletal Muscle. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.