

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB03004 - Goat Anti-phospho-CD244 / 2B4 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** CD244, CD244 molecule, natural killer cell receptor 2B4, 2B4, NAIL, NKR2B4, Nmrk, SLAMF4, CD244 natural killer cell receptor 2B4, NK cell activation

inducing ligand NAIL, natural killer cell receptor 2B4

Official Symbol: CD244

Accession Number(s): NP\_057466.1

Human GeneID(s): 51744

#### **Immunogen**

Peptide with sequence C-EFLTIpYEDVKD, from the internal region of the protein sequence according to NP\_057466.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:64000.

Immunoprecipitation: Approx 75kDa band observed in Human Natural Killer (NK) cell lysates after with immunoprecipitation mouse monoclonal antibody C1.7 coupled to Protein G agarose (calculated MW of 41.1kDa according to NP\_057466.1). We do note that CD244 is a glycoprotein which makes a higher size band to be expected. Recommended concentration: 1-3µg/ml.

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB03004 (2µg/ml) staining of IP from a lysate of Human Natural Killer cells with (+) or without (-) pervanadate treatment of the cells. Primary incubation was 1 hour. Detected by chemiluminescence. Data kindly provided by Dr Nicholas G Clarkson, Sir William Dunn School of Pathology, Oxford, UK