



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06437 - Goat Anti-Pirin Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: pirin, PIR, pirin (iron-binding nuclear protein)

Official Symbol: PIR

Accession Number(s): NP_003653.1

Human GeneID(s): [8544](#)

Important Comments: Variants (NP_003653.1; NP_001018119.1) encode the same protein

Immunogen

Peptide with sequence GSSKKVTLVSLSREC, from the N Terminus of the protein sequence according to NP_003653.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.


Western blot: Approx 33kDa band observed in Human Heart lysate (calculated MW of 32.1kDa according to NP_003653.1). Recommended concentration: 0.3 - 1.0 µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Liver and Spleen. Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human, Mouse



EB06437 (0.3 μ g/ml) staining of Human Heart lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB06437 (5 μ g/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB06437 (5µg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.