

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06239 - Goat Anti-PIST / FIG / GOPC Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** PIST, GOPC, CAL, PDZ/coiled-coil domain binding partner for the rho-family GTPase TC10, fused in glioblastoma, CFTR-associated ligand, Golgi associated PDZ and coiled-coil motif containing protein, golgi associated PDZ and coiled-coil motif containing, CAL, FIG, GOPC1, dJ94G16.2, dJ94G16.2 PIST

Official Symbol: GOPC

Accession Number(s): NP\_065132.1; NP\_001017408.1

Human GeneID(s): 57120

Important Comments: This antibody is expected to recognize both reported isoforms

(NP\_065132.1; NP\_001017408.1).

## **Immunogen**

Peptide with sequence C-LDDLHTLYHKKSY, from the C Terminus of the protein sequence according to NP\_065132.1; NP\_001017408.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx. 55kDa band observed in Human Brain (Frontal Cortex) lysates (calculated MW of 50.5kDa according to NP\_065132.1). Recommended concentration: 0.05-0.2µg/ml.

Immunofluorescence: This product has been successfully used on HeLa cells.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Dog, Cow



EB06239 (0.05μg/ml) staining of Human Frontal Cortex lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.